



Sheena Black, MD

Orthopaedic Surgery, Sports Medicine 469-800-7200

Triceps Tendon Repair

<u>Postop</u>	Range of Motion	<u>Immobilizer</u>	<u>Exercises</u>
Phase I Weeks 0-2 PT 1-2x/week	None	Splint at all times at 30-45° flexion	Hand/wrist ROM as tolerated Limit shoulder flexion to 90° Gentle shoulder PROM NO weight bearing activities (pushing open door, pushing up from a chair)
Phase II Weeks 3-6 PT 1-2x/week HEP daily	Elbow ROM/brace: Week 3: 0 – 45° Week 4: 0 – 60° Week 5-6: 0 – 90°	Brace at ALL times (except for exercise and bathing)	Continue shoulder/wrist/hand ROM and isometrics Active assist pronation/supination at position of max flexion NO active elbow extension x 6 weeks
Phase III Weeks 6-8 PT 2-3x/week HEP daily	Week 7-8: 0 – 125°	Brace at ALL times (except for exercise and bathing)	Progress AROM of shoulder Initiate active, concentric elbow extension (no resistance) in pain-free range NO eccentric triceps exercises
Phase IV Weeks 8-12 PT 2-3x/week HEP daily	Full	Discontinue brace if good motor control	Initiate sub-maximal triceps isometrics Eccentric triceps activity (no resistance) Progress biceps strengthening Resisted SA punch & bear hugs Resisted IR and ER at 30° Abduction progress to 90° Prone scapular series OK for stationary bike and light jogging
Phase V Weeks 12-16 PT 2-3x/week HEP daily	Full	None	Progress triceps strengthening Bicep curls with dumbbells IR/ER exercises at 90° abduction CKC UE weight-bearing: wall weight shifts, quadruped rocking Initiate plyotoss (double arm – progress to single arm) Initiate pushup progression at week 14 (limit elbow flexion to 45 degrees to start) Interval throwing program at week 16-20 Initiate light upper body plyometric program at week 16-20 PNF/ Diagonal pattern strengthening Initial sport specific activities