



Sheena Black, MD

Orthopaedic Surgery, Sports Medicine DrBlack@SheenaBlackMD.com

PHYSICAL THERAPY PRESCRIPTION MENISCAL REPAIR – ALL-INSIDE

Name:			DOB:					
Post-Operative	e Diagnosis:	□ Right	□ Left					
Additional Pro	cedures:	□ Latera	l Meniscal Repair	Medial Meniscal Repair				
		□ Latera	al Menisectomy	Medial Menisectomy				
		□ Other:						
Surgical Date:								
Weight Bearing: 🛛 WBAT								
Weight bearing modifications:								
ROM: Full passive knee extension, <u>limit knee flexion to 0 – 90 degrees</u> ROM Modifications:								
Brace Instructions:								
Additional Instructions:								
Frequency & Duration: 1-2 times per week for 6 - 8 weeks								
Rx:	Dressing change / wound check 2-3 days post-op							
	Call my office if patient unable to get $0 - 90^{\circ}$ ROM by 2 weeks post-op							
	Emphasize early full passive extension to 0° (No recurvatum for hamstring grafts)							
Precautions: X Per my PT protocol Other:								
Physician's Signature:,MD								





Sheena Black, MD

Orthopaedic Surgery, Sports Medicine DrBlack@SheenaBlackMD.com

PHYSICAL THERAPY PROTOCOL KNEE ARTHROSCOPY – MENISCAL REPAIR – ALL-INSIDE

	<u>Weight</u> <u>Bearing</u>	<u>Brace</u>	<u>ROM</u>	<u>Exercises</u>
Phase I Weeks 0-2 PT 1-2x/week HEP daily	WBAT in brace locked in full extension w/ crutches *	Locked in full extension for sleeping and all activity Off for exercises & hygiene	0-90° when non- weight bearing	Heel slides, quad sets, patella mobs SLR No weight bearing with flexion > 90°
Phase II Weeks 2-6 PT 2-3x/ week HEP daily	 2-4 weeks: Full in brace unlocked 0-90° 4-6 weeks: Full w/o brace 	2-6 weeks : Unlocked 0-90° Off at night Discontinue brace at 6 weeks	As tolerated	Continue Phase 1 Heel raises, closed chain, terminal knee extension avoiding tibial rotation Activities w/ brace until 4-6 weeks After 6 weeks, w/o brace as tolerated No weight bearing with flexion > 90 °
Phase III Weeks 8-12 PT 2-3x/week HEP daily	WBAT	None	Full	Progress closed chain activities Begin hamstring work, lunges/leg press 0-90°, proprioception exercises, balance/core/hip/glutes Begin stationary bike
Phase IV Weeks 12-20 PT 1-2x/week HEP daily	WBAT	None	Full	Progress Phase III exercises & functional activities Single leg balance, core, glutes, eccentric hamstrings, elliptical, bike Ok to progress to swimming at 12 weeks Advance to sport-specific drills & running/jumping at 16 weeks once cleared by MD

IF QUESTIONS: PLEASE CALL 469-800-7200